

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C08F 8/30, C12N 11/08	A1	(11) International Publication Number: WO 96/26223 (43) International Publication Date: 29 August 1996 (29.08.96)
--	----	---

(21) International Application Number: PCT/GB96/00428
(22) International Filing Date: 26 February 1996 (26.02.96)
(30) Priority Data:
9503749.5 24 February 1995 (24.02.95) GB

(81) Designated States: JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).



Published
With international search report.

(71) Applicant (for all designated States except US): BRITISH BIOTECH PHARMACEUTICALS LIMITED [GB/GB]; Watlington Road, Cowley, Oxford OX4 5LY (GB).

(72) Inventors; and
(75) Inventors/Applicants (for US only): FLOYD, Christopher, David [GB/GB]; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB). LEWIS, Christopher, Norman [GB/GB]; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB).

(74) Agent: WALLS, Alan, J.; British Biotech Pharmaceuticals Limited, Watlington Road, Cowley, Oxford OX4 5LY (GB).

(54) Title: SYNTHESIS OF HYDROXAMIC ACID DERIVATIVES



(57) Abstract

A solid phase reaction component comprising a solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of covalently bound hydroxylamine or protected hydroxylamine groups of formula (A) or (B), wherein P_1 is hydrogen or an amino protecting group and P_2 is hydrogen or a hydroxyl protecting group, and the bond designated (a) is one which covalently links the group (A) or (B) to the residue of the solid substrate and is cleavable under acid conditions or by photolysis. Such components are useful in the solid phase synthesis of, for example, compounds which are matrix metalloproteinase inhibitors.

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Synthesis of Hydroxamic Acid Derivatives

The present invention relates to a solid phase reaction component, and to a process for the use of such a component in the synthesis of individual hydroxamic acid derivatives or a combinatorial library of such compounds. In particular, the solid phase reaction component may be used for the synthesis of hydroxamic acids that are inhibitors of zinc metalloproteinase enzymes. Such enzymes are involved in tissue degradation and the release of tumour necrosis factor from cells.

Background to the Invention

Solid Phase Synthesis

Solid phase synthesis is an established and effective method for the preparation of peptides, and offers advantages over conventional solution phase chemistry in terms of purification and simplicity (Atherton E, Sheppard RC, *Solid Phase Peptide Synthesis: A Practical Approach*; IRL Press at Oxford University Press: Oxford, 1989). Solid phase synthesis may also be used for the preparation of non-peptide molecules (Leznoff CC, *Acc. Chem. Res.*, 1978, 11, 327-333) and recently there has been considerable interest in the application of this methodology to the synthesis of combinatorial libraries for biologically active lead compound optimisation and discovery (Moos WH *et al.*, *Annu. Rep. Med. Chem.*, 1993, 28, 315-324).

Solid phase synthesis requires an appropriate solid substrate which carries a plurality of functional groups to which the first reactive entity in the proposed synthesis may be covalently coupled, and from which the desired molecule may be cleaved after assembly. The solid substrate should be compatible with the solvents and reaction conditions that are to be used in the peptide or non-peptide synthesis.

The final step in solid phase synthesis is the cleavage of the covalent bond between the desired peptide or non-peptide molecule and the linker. It is desirable that the conditions for the cleavage are orthogonal to those used during the reactions

employed for the synthesis of the peptide or non-peptide on the solid support such that inadvertent cleavage does not occur during the synthesis. Furthermore, the conditions for cleavage should be relatively mild such that they do not result in degradation of the desired peptide or non-peptide. Solid substrates which present hydroxyl groups as the points of attachment for the first stage of the synthesis are commonly used, for example substrates which present hydroxyl groups as derivatives of benzyl alcohol, the peptide or non-peptide being attached as a benzyl ester and cleaved by hydrolysis, transesterification or aminolysis to release the peptide or non-peptide as a carboxylic acid, carboxylate ester or as a carboxamide. Also used are substrates which present amino groups, for example as derivatives of diphenylmethanamine, the peptide or non-peptide being attached as a carboxamide and cleaved by hydrolysis to release the peptide or non-peptide as a carboxamide. Substitution of such linkers by a nitro group can enable the photolytic cleavage of the peptides or non-peptides from the residue of the solid substrate.

Hydroxamic Acid Derivatives

Certain hydroxamic acid derivatives possess useful biological activities. Examples of such hydroxamic acids include compounds that inhibit urease (Otake S *et al.*, *Chem. Pharm. Bull.*, 1992, **40**, 2764-2768), trypanosome glycerol-3-phosphate oxidase (Grady *et al.*, *Mol. Biochem. Parasitol.*, 1986, **19**, 231-240), dehydropeptidase-1 (EP-B-276,947), ribonucleotide reductase (Farr RA *et al.*, *J. Med. Chem.*, 1989, **32**, 1879-1885), 5-lipoxygenase (Kerdesky FAJ *et al.*, *Tetrahedron Lett.*, 1985, **26**, 2134-2146; US-4,731,382), substance P degradation (Laufer R *et al.*, *Eur. J. Biochem.*, 1985, **150**, 135-140), cardiovascular metalloproteinase enzymes (Turbanti L *et al.*, *J. Med. Chem.*, 1993, **36**, 699-707; WO-9428012) and matrix metalloproteinase enzymes (Schwartz MA, Van Wart HE, *Prog. Med. Chem.*, 1992, **29**, 271-334).

Compounds which have the property of inhibiting the action of metalloproteinases involved in connective tissue breakdown such as collagenase, stromelysin and gelatinase (known as "matrix metalloproteinases", and herein referred to as MMPs)

are thought to be potentially useful for the treatment or prophylaxis of conditions involving such tissue breakdown, for example rheumatoid arthritis, osteoarthritis, osteopenias such as osteoporosis, periodontitis, gingivitis, corneal, epidermal or gastric ulceration, and tumour metastasis, invasion and growth. It has been found that hydroxamic acid MMP inhibitors can also inhibit the production of the cytokine tumour necrosis factor (herein referred to as "TNF") (Mohler *et al.*, *Nature*, 1994, 370, 218-220; Gearing AJH *et al.*, *Nature* 1994, 370, 555-557; McGeehan GM *et al.*, *Nature* 1994, 370, 558-561). Compounds which inhibit the production or action of TNF are thought to be potentially useful for the treatment or prophylaxis of many inflammatory, infectious, immunological or malignant diseases. These include, but are not restricted to, septic shock, haemodynamic shock and sepsis syndrome, post ischaemic reperfusion injury, malaria, Crohn's disease, mycobacterial infection, meningitis, psoriasis, congestive heart failure, fibrotic disease, cachexia, graft rejection, cancer, autoimmune disease, rheumatoid arthritis, multiple sclerosis, radiation damage, toxicity following administration of immunosuppressive monoclonal antibodies such as OKT3 or CAMPATH-1 and hyperoxic alveolar injury. Since excessive TNF production has been noted in several diseases or conditions also characterised by MMP-mediated tissue degradation, compounds which inhibit both MMPs and TNF production may have particular advantages in the treatment or prophylaxis of diseases or conditions in which both mechanisms are involved.

Classes of MMP Inhibitors

The known hydroxamic acid MMP inhibitors may be grouped into three classes; i) peptidyl hydroxamates, ii) succinyl hydroxamates and iii) arylsulfonamido hydroxamates.

- i) The following patent publication discloses peptidyl hydroxamic acid-based MMP inhibitors:

EP-A-345359 (Fuji)

The tri- and tetra-peptidyl hydroxamic acid derivatives disclosed in the above publication and described elsewhere (Otake S *et al.*, *Chem. Pharm. Bull.*, 1990, **38**, 1007-1011; Otake S *et al.*, *Chem. Pharm. Bull.*, 1991, **39**, 1489-1494; Otake S *et al.*, *Biochem. Biophys. Res. Commun.*, 1994, **199**, 1442-1446) can be regarded as having the following basic structure (I):




wherein the four groups R_1 , X_1 , X_2 and X_3 may vary according to the detailed disclosure of each of the publications.

ii) The following patent publications disclose succinyl hydroxamic acid-based MMP inhibitors:

US 4599361	(Searle)
EP-A-0236872	(Roche)
EP-A-0274453	(Bellon)
WO 90/05716	(British Bio-technology)
WO 90/05719	(British Bio-technology)
WO 91/02716	(British Bio-technology)
EP-A-0489577	(Celltech)
EP-A-0489579	(Celltech)
EP-A-0497192	(Roche)
WO 92/13831	(British Bio-technology)
WO 92/22523	(Research Corporation Technologies)
WO 93/09090	(Yamanouchi)
WO 93/09097	(Sankyo)
WO 93/20047	(British Bio-technology)
WO 93/24449	(Celltech)
WO 93/24475	(Celltech)
US 5256657	(Sterling Winthrop)
EP-A-0574758	(Roche)

WO 94/02446	(British Bio-technology)
WO 94/02447	(British Bio-technology)
WO 94/21612	(Otsuka)
WO 94/25434	(Celltech)
WO 94/25435	(Celltech)

The hydroxamic acid derivatives disclosed in the above publications can be regarded as having the following basic structure (II):




wherein the five substituents $R_2 - R_6$ may vary according to the detailed disclosure of each publication. The balance of intrinsic level of activity, degree of specificity of inhibition of particular categories of MMP, physicochemical and pharmacokinetic properties can vary in an unpredictable way as the substituents $R_2 - R_6$ are varied.

iii) The following patent publication discloses arylsulfonamido hydroxamic acid-based MMP inhibitors:

EP-A-606046 (Ciba-Geigy)

The hydroxamic acid derivatives disclosed in the above publication can be regarded as having the following basic structure (III):



wherein the four substituents R_1 - R_4 and Ar may vary according to the detailed disclosure of the publication.


Brief Description of the Invention

A key step in the synthesis of many hydroxamic acid derivatives is the reaction of a carboxylic acid group with hydroxylamine or a protected hydroxylamine by first activating the carboxylic acid or by conducting the reaction in the presence of an activating agent. In a recent patent application the possibility that peptidyl hydroxamic acids might be prepared using solid phase synthesis was recognised (WO 94/28012). However, the proposed protocol involved first synthesis of the peptide on a solid support followed by cleavage from the support and subsequent conversion of the C-terminal carboxylic acid group to a hydroxamic acid by a method analogous to that indicated above.


It was the hypothesis of the present inventors that hydroxamic acid derivatives might be prepared by the reaction of a solid substrate presenting hydroxylamine groups as the point of attachment for the first reactive entity of the synthesis, then reacting the resultant solid phase intermediate with the further reactive entities required for the desired synthesis, followed by a cleavage step to release the final hydroxamic acid derivative. Such a hydroxylamine-presenting solid phase reaction component could simplify the synthesis and purification of biologically active hydroxamic acid derivatives, for example MMP inhibitors of the three classes referred to above, and could be applicable to the automation of such syntheses. In addition it would enable the solid phase synthesis of combinatorial libraries of hydroxamic acids. Thus, this invention makes available such hydroxylamine-presenting solid phase reaction components, and provides a process for their use in solid phase synthesis of hydroxamic acid derivatives.

Detailed Description of the Invention

The present invention provides, in one of its aspects, a solid phase reaction component comprising a solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of covalently bound hydroxylamine or protected hydroxylamine groups of formula (A) or (B)



(A)




(B)

wherein P_1 is hydrogen or an amino protecting group and P_2 is hydrogen or a hydroxyl protecting group, and the bond designated (a) is one which covalently links the group (A) or (B) to the residue of the solid substrate and is cleavable under acid conditions or by photolysis.

The bond designated (a) covalently links the group (A) or (B) to the residue of the solid substrate. The residue of the solid substrate will usually comprise a base substrate carrying suitable linker groups which indirectly link the hydroxylamine group (A) or (B) to the base substrate. Suitable base substrates include those known in the art of solid phase peptide synthesis (see for example those described in Stewart JM and Young JD, *Solid Phase Peptide Synthesis*, 2nd Ed; Pierce Chemical Company: Rockford, Illinois, 1984). They include inorganic substrates, for example kieselguhr, silica gel and controlled pore glass, and polymeric organic substrates, for example polystyrene, polypropylene, polyethylene glycol, polyacrylamide, cellulose, as well as composite inorganic/polymeric substrates such as polyacrylamide supported within a matrix of kieselguhr particles. Such known base substrates include amino and hydroxy functionalised solid substrates, ie those which are chemically modified by introduction of amino or hydroxyl groups, to serve as


convenient points for further chemical manipulation. Examples of particular amino or hydroxy functionalised base substrates are:

hydroxymethyl copoly(styrene-1 or 2% divinylbenzene), which can be represented as



wherein P represents the polymer backbone;

benzhydramine copoly(styrene-1 or 2% divinylbenzene) ("BHA Resin") or methyl benzhydramine copoly(styrene-1 or 2% divinylbenzene) ("MBHA Resin"), which can be represented as



wherein P represents the polymer backbone and R is hydrogen or methyl respectively;

polyethylene glycol polystyrene ("PEG-PS");

poly(dimethylacrylamide)polystyrene composite ("Polyhipe");


polyacrylamide Kieselguhr composite ("Macrosorb");

or functionalised controlled pore glass.

Thus, in the case of peptide synthesis, hydroxyl- or amino-carrying linker groups can be introduced onto such amino and hydroxy functionalised base substrates. In case of a hydroxyl-carrying linker group, the first amino acid of the peptide to be constructed may be attached as an ester formed between the linker-presented hydroxyl group and the carboxyl group of the amino acid. In the case of amino-carrying linker groups, the first amino acid of the peptide to be constructed may be

attached as a carboxamide formed between the linker-presented amino group and the carboxyl group of the amino acid.

In an analogous fashion, the solid phase reaction components of the present invention may comprise a base substrate, for example of the kind referred to above, and a linker group which presents the hydroxylamine group (A) or (B) for reaction with the first reactive entity in the proposed synthesis. A preferred embodiment of this aspect of the invention is a solid phase reaction component, comprising a solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of groups of formula (IV):



wherein

R_{10} represents hydrogen or $(phenyl)CH_2-$, optionally substituted in the phenyl ring by one or more substituents selected from C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, nitrile or NO_2 ;

R_{11} represents hydrogen, C_1-C_6 alkyl, or phenyl optionally substituted by one or more substituents selected from C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, nitrile or NO_2 ;


R_{12} and R_{12A} independently represent hydrogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, nitrile or NO_2 ;

R_{13} represents a group $-(X')_q-Y-$ wherein q is 0 or 1, X' represents $-C(=O)-$,

-CH₂-, -CH₂C(=O)-, -O(CH₂)_nC(=O)-, -O(CH₂)_nC(=O)-(A')_m-, or
 -O(CH₂)_nC(=O)-(A')_m-B'-, wherein n is an integer from 1 to 6, m is 0 or 1, A'
 represents -O-CH(R')-NH- wherein R' is the side chain of a natural or
 unnatural alpha amino acid, B' represents a spacer group -NH(CH₂)_p- wherein
 p is 0 or an integer from 1 to 6, and Y represents -O- or -NH-.

In this embodiment, the group (IV) is linked to the solid substrate via "Y", defined as -
 O- or -NH-. It will be apparent that these "Y" groups may be incorporated during
 synthesis of the solid phase reaction component of the invention, by starting with an
 amino or hydroxy functionalised base substrate, for example
 hydroxymethylpolystyrene, hydroxymethyl copoly(styrene-1%divinylbenzene),
 benzhydrylaminepolystyrene, benzhydrylamine copoly(styrene-1%divinylbenzene
 ("BHA Resin"), methyl benzhydrylaminepolystyrene, methyl benzhydrylamine
 copoly(styrene-1%divinylbenzene, ("MBHA Resin"); polyethylene glycol polystyrene
 ("PEG-PS"); poly(dimethylacrylamide)polystyrene composite ("Polyhipe");
 polyacrylamide Kieselguhr composite ("Macrosorb"); or functionalised controlled
 pore glass.

Another preferred embodiment of this aspect of the invention is a solid phase
 reaction component, comprising a solid substrate, substantially insoluble in aqueous
 or organic reaction media, carrying a plurality of groups of formula (IVA):



wherein R₁₀, R₁₁, R₁₂ and R_{12A} are as defined in formula (IV), R_{11A} is as defined for R₁₁
 in formula (IV), and R_{13A} is a bond or is as defined for R₁₃ in formula (IV).

As used herein the term "C₁-C₆ alkyl" means a straight or branched chain alkyl
 moiety having from 1 to 6 carbon atoms, including for example, methyl, ethyl, n-

propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, pentyl and hexyl. "C₁-C₆ alkoxy" means an alkoxy group wherein the alkyl part is C₁-C₆ alkyl.

In the preferred solid phase reaction component of the invention carrying a plurality of groups of formula (IV), referred to above:

R₁₀ may for example be hydrogen, 4-methoxybenzyl or 2,4-dimethoxybenzyl. Presently preferred are compounds in which R₁₀ is hydrogen.

R₁₁ may for example be hydrogen, methyl, phenyl, 4-methylphenyl, 4-methoxyphenyl or 2,4-dimethoxyphenyl. Presently preferred are compounds in which R₁₁ is hydrogen.

R₁₂ may for example be hydrogen or methoxy. Presently preferred are compounds in which R₁₂ is hydrogen.

R₁₃ together with the base substrate to which it is attached may for example be oxymethyl-copoly(styrene-1 divinylbenzene)resin, oxymethyl-copoly(styrene-2% divinylbenzene)resin, oxyacetomidomethylpolyethyleneglycol-copoly(styrene-1% divinylbenzene)resin or oxyacetomidomethylpolyethyleneglycol-copoly(styrene-2% divinylbenzene)resin. Presently preferred are compounds in which R₁₃ together with the base substrate to which it is attached is oxymethyl-copoly(styrene-1% divinylbenzene)resin.

Specific preferred solid phase reaction components of the invention carrying a plurality of groups of formula (IV), are:

4-(O-Methylhydroxylamine)phenoxy-methyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh),


4-[4-(O-Methylhydroxylamine)-3-methoxyphenoxy]-(N-4-methylbenzhydryl)butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh),

4-(2',4'-Dimethoxyphenyl-O-methylhydroxylamine)-phenoxymethyl-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh), and

4-[4-(1-Aminooxyethyl)-2-methoxy-5-nitrophenoxy]-(N-4-methylbenzhydryl)butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh).

A specific preferred solid phase reaction component of the invention carrying a plurality of groups of formula (IVA), is: O-Hydroxylamine-2'-chlorotrityl-copoly(styrene-1%-divinylbenzene)-resin (200-400 mesh)

Solid phase reaction components of the invention may be prepared by standard synthetic techniques from solid substrates which are in general commercially available or readily derivable from commercially available materials. For example, the preferred solid phase reaction component of the invention carrying a plurality of groups of formula (IV), referred to above may be prepared by treating a solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of benzyl alcohol groups of general formula (V).




wherein R_1 , R_{13} are as defined for general formula (IV) with a compound of formula (VI)



(VI)


wherein R_{10} is as defined for general formula (IV) but is other than hydrogen under Mitsunobu reaction conditions using triphenylphosphine and diethylazodicarboxylate or similar reagents. The preferred solid phase reaction component of the invention carrying a plurality of groups of formula (IV), referred to above, wherein R_{10} is hydrogen may be prepared by treating a solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of groups of general formula (VII)



(VII)

wherein R_{11} - R_{13} are as defined for general formula (IV) with hydrazine.

A solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of groups of general formula (VII) can be prepared by coupling a phthalimide derivative of formula (VIII) with a solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of benzyl alcohol groups of general formula (V) defined above



(VIII)

wherein R_{11} - R_{13} are as defined in general formula (IV) under Mitsunobu reaction conditions using triphenylphosphine and diethyl azodicarboxylate or similar reagents. Solid substrates, substantially insoluble in aqueous or organic reaction media, carrying a plurality of groups of general formula (V), and compounds of formulae (VI) and (VIII), are known in the art or may be prepared by procedures known to those skilled in the art.

In an analogous manner, the preferred solid phase reaction components of formula (IVA) above may be prepared. Alternatively, the hydroxylamine group may be introduced by displacement of a leaving group present in the base substrate, eg a halogen group such as chloro, using a suitably protected hydroxylamine derivative, eg a compound of formula (Viii) above.



The choice of solvent for syntheses based on a solid phase reaction component of the invention will of course depend on the nature of the reagents to be reacted with such component, but will also be influenced by the nature of that component. For example many of the polymer reaction components of the invention may swell to a greater or lesser extent in certain solvents, and generally such swelling will be desirable for the efficiency of the reaction with other reagents in the desired synthesis.

Solid phase reaction components of the invention presenting groups of type (B) above will generally be accessible (i) by displacement of a leaving group (eg a triflate, mesylate or halogen group) from the desired base substrate carrying such leaving groups, using hydroxylamine or O-protected hydroxylamine, or (ii) by reacting the desired base substrate carrying carbonyl groups with hydroxylamine or O-protected hydroxylamine to form oxime groups, and then reducing the oxime double bond, eg using a metal hydride.

In another of its aspects, the present invention comprises a process for the preparation of a desired synthetic product whose structure is characterised by the

presence of a covalently bonded hydroxamic acid group -CONHOH, which process comprises the steps of:

- (i) forming a mixture of a liquid reaction medium and a solid phase reaction component which is substantially insoluble in the said liquid reaction medium and carries a plurality of covalently bound moieties of formula (A1) or (B1)



in which formulae X represents the residual, non-hydroxamate, partial structure of the desired product, P₁ represents hydrogen or an amino-protecting group, P₂ represents hydrogen or a hydroxyl protecting group, and the bond designated (a) is one which covalently links the moieties (A1) or (B1) to the residue of the solid substrate and is cleavable under acid conditions or by photolysis; and

- (ii) in the resultant mixture, cleaving the said bond designated (a) and, if P₁ or P₂ as the case may be is not hydrogen, removing that protecting group P₁ or P₂ before, after or during cleavage of bond (a); and

- (iii) separating the resultant liquid reaction phase from the resultant reaction solids and recovering the desired product from the separated liquid reaction phase.


Trifluoroacetic acid will generally be suitable for the acid hydrolysis of the bond (a). Depending on the structure of the linker group, solutions of trifluoroacetic acid ranging from 95%v/v to 1%v/v may be used. In Example 5 below, the linker group is photolytically cleavable, and in the remaining Examples it is cleavable by acid hydrolysis.


In step (i) of the above process the solid phase reaction component which carries a plurality of moieties of formula (A1) or (B1) may be derived by appropriate chemical modification from the solid phase reaction component of the invention, described above, carrying a plurality of hydroxylamine or protected hydroxylamine groups of formula (A) or (B).

In the process of the invention, the solid phase reaction component may be in the form of a finely divided solid, or a web, membrane or open pore matrix. For example, the liquid reaction medium may be mixed with the solid phase reaction product by passing the former through a liquid permeable bed or column of the latter, or by contacting the former with a discrete quantity of the latter in finely divided form in a reaction vessel.

The solid phase reaction component and process of the present invention are applicable in syntheses of the following general types, Schemes 1 and 2:

Scheme 1





In schemes 1 and 2, the vertical wavy line represents the base solid substrate, L represents a linker group, P₁ and P₂ are as defined above, and B₁...B_n are the reaction building blocks for the synthesis of the desired product. In both cases, the desired product is released from the solid phase support by acid hydrolysis or photolysis of the bond designated (a) in formula (A), or (B), and the protecting group P₁ or P₂ may be removed in the same or a separate step depending on its identity.

If desired, the intermediate solid phase reaction product can be isolated and physically partitioned into a plurality of portions after the addition of each building block B₁...B_n. Each resultant portion may then be reacted with a different next building block. This process facilitates the parallel synthesis of a multiplicity of different hydroxamate end products.

Alternatively, the solid phase reaction component can be physically partitioned into a plurality of portions and each resultant portion may then be reacted with a different first building block B₁. The separate portions may then be mixed to form a single portion which is then split into a plurality of portions. Each resultant portion may then be reacted with a different second building block. The separate portions may then be mixed to form a single portion which is then split into a plurality of portions for the coupling of the different second building blocks B₂. This process may be


repeated a number of times until the final building block B_n has been added. This process facilitates the combinatorial synthesis of a mixture of a multiplicity of different hydroxamate end products.

Specific applications of the process of the invention are the preparation of hydroxamic acids of the three classes referred to above, namely peptidyl, succinyl and arylsulphonamido hydroxamates.

For the synthesis of peptidyl hydroxamates (formula (I) above) by, for example, Scheme 1, the building blocks $B_1 \dots B_n$ represent the amino acids for sequential coupling. The first amino acid is coupled to the groups A carried by solid phase reaction component of the invention, for example one of the preferred solid phase reaction components of the invention carrying a plurality of groups of formula (IV) or (IVA). In the latter case, the amino group of the amino acid component may be protected as a 9-fluorenylmethoxycarbonyl ("Fmoc") derivative and the coupling of the amino acid carboxyl group to the nitrogen of the group A carried by the solid phase reaction component of the invention may be facilitated by the use of a coupling agent such as is commonly used in peptide synthesis, for example a carbodiimide (e.g. diisopropylcarbodiimide), a phosphonium salt (e.g. benzotriazol-1-yl-oxy-tris-(dimethylamino)phosphonium hexafluorophosphate) or a uronium salt (e.g. 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate), with the optional addition of an additive such as 1-hydroxybenzotriazole. Alternatively, the coupling reaction may be conducted by preforming an active ester derivative (e.g. pentafluorophenyl ester) of the carboxylic acid component and optionally conducting the reaction in the presence of an additive such as 1-hydroxybenzotriazole. The Fmoc amino protecting group may then be removed by treatment of the reaction product with a basic amine such as piperidine. During the coupling reaction any groups in the amino acid which are potentially reactive under the coupling conditions may be protected from such reaction as is usual in peptide synthesis. Addition of the second and any subsequent amino acids proceeds in a similar manner until the desired peptidyl hydroxamate is finally cleaved from the

solid substrate, eg by acid hydrolysis of the bond designated (a), and any protecting groups present in the molecule may be removed before during or after such cleavage, depending on the nature of the protecting groups in question.


For the synthesis of succinyl hydroxamates (formula (II) above) by, for example, Scheme 1, an appropriate choice of building blocks B_1, \dots, B_n to effect the synthesis is made. The first building block will generally be a carboxylic acid (also containing a site suitable for coupling the next building block) which is coupled to the groups A carried by solid phase reaction component of the invention, for example one of the preferred solid phase reaction components of the invention carrying a plurality of groups of formula (IV) or (IVA). Again the coupling of the carboxyl group of the first building block to the nitrogen of the group A carried by the solid phase reaction component of the invention may be facilitated by the use of a coupling agent such as is commonly used in peptide synthesis, for example a carbodiimide (e.g. diisopropylcarbodiimide), a phosphonium salt (e.g. benzotriazol-1-yl-oxy-tris-(dimethylamino)phosphonium hexafluorophosphate), or a uronium salt (e.g. 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate), with the optional addition of an additive such as 1-hydroxybenzotriazole. Alternatively, an activated ester, for example a pentafluorophenyl ester, of the first carboxylic acid building block may be coupled to the group A carried by the solid phase reaction component of the invention in the presence of dimethylaminopyridine. A particular first carboxylic acid building block may be one of formula (IX)



(IX)

wherein R_2 and R_3 are groups dictated by the desired succinyl hydroxamate final product and R_1 is a carboxyl protecting group which is compatible with solid phase synthesis such as an allyl group. Where the compound (IX) is coupled to the solid

phase reaction components of the invention in this way, the $R_{1,4}$ carboxyl protecting group in the resultant product may be converted to a carboxylic acid group by deprotection. In the case where $R_{1,4}$ is an allyl group deprotection may be achieved by palladium catalysis. Suitable reagents for this step are $\text{Pd}(\text{PPh}_3)_4$ or $\text{Pd}(\text{OAc})_2$ / triphenylphosphine, and the reaction is usually conducted in the presence of an allyl acceptor such as morpholine. The carboxylic acid group thus generated may then be used as the point of attachment for the second building block, which will generally be an amine of formula (X)




(X)

wherein R_4 - R_6 are again groups dictated by the desired succinyl hydroxamate final product. The methods described above and common in peptide synthesis are used to effect the coupling of the amine group of (X) with the carboxylic acid group of the first building block now carried on the solid phase reaction component. During the coupling reactions of the foregoing assembly process, any groups in the first and second building blocks which are potentially reactive under the coupling conditions may be protected from such reaction as is usual in peptide synthesis. After assembly of the desired succinyl hydroxamate, the product may be cleaved from the solid substrate, eg by acid hydrolysis of the bond designated (a), and any protecting groups present in the molecule may be removed before, during or after such cleavage, depending on the nature of the protecting groups in question.

For the synthesis of arylsulphonamide hydroxamates (formula (III) above) by the method of, for example, Scheme 1, an appropriate choice of building blocks $B_1 \dots B_n$ to effect the synthesis is made. The first building block will either be an alpha amino acid in which the alpha amino group is protected (for example as a 9-fluorenylmethoxycarbonyl ("Fmoc") or allyloxycarbonyl ("Aloc") derivative), and the

substituents on the alpha C atom (corresponding to R_7 and R_8 in formula (III)) are dictated by the desired arylsulphonamide hydroxamate final product or an alpha halo acid of formula (XI)



wherein R_7 and R_8 are dictated by the desired arylsulphonamide hydroxamate final product and Hal is a halogen such as bromo. This first building block (protected alpha amino acid or alpha halo acid) is coupled to the groups A carried by solid phase reaction component of the invention, for example one of the preferred solid phase reaction components of the invention carrying a plurality of groups of formula (IV) or (IVA). In the latter case, the coupling of the protected alpha-amino acid or alpha-halo acid carboxyl group to the nitrogen of the group A carried by the solid phase reaction component of the invention may be facilitated by the use of a coupling agent such as is commonly used in peptide synthesis, for example a carbodiimide (e.g. diisopropylcarbodiimide), a phosphonium salt (e.g. benzotriazol-1-yl-oxy-tris-(dimethylamino)phosphonium hexafluorophosphate) or a uronium salt (e.g. 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate), with the optional addition of an additive such as 1-hydroxybenzotriazole. Alternatively this first building block may be coupled to the group A carried by the solid phase reaction component of the invention as an activated ester, for example a pentafluorophenyl ester, in the presence of dimethylaminopyridine.

In the case where the first building block is a protected alpha-amino acid the N-protecting group (eg Fmoc or Alloc) may be removed by treatment of the reaction product with a basic amine such as piperidine in the case of base labile protecting groups (e.g. Fmoc) or treatment with a suitable palladium catalyst in the case of palladium labile protecting groups (e.g. Alloc). The free amino group then serves as

the site for formation of the appropriate arylsulphonamide and alkylation of the sulphonamide nitrogen to complete the synthesis of the desired arylsulphonamide hydroxamate, which can then be cleaved from the solid substrate, eg by acid hydrolysis of the bond designated (a).

In the case where the first building block is an alpha-halo acid the halogen may be displaced by reaction with an amine to form a secondary amine which serves as the site for formation of the sulphonamide to complete the synthesis of the desired arylsulphonamide hydroxamate, which can then be cleaved from the solid substrate, eg by acid hydrolysis of the bond designated (a).

Examples 1-5 disclose the preparation of solid phase reaction components of the invention and Examples A-H illustrate their utility in the preparation of biologically active hydroxamic acid derivatives.

The amino acids used in the examples below were commercially available or were prepared according to literature procedures.

The following abbreviations have been used throughout:


BPB	Bromophenol blue
DMF	N,N-Dimethylformamide
DMSO	Dimethylsulfoxide
Fmoc	9-Fluorenylmethoxycarbonyl
pyBOP	Benzotriazol-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate
HOBt	N-Hydroxybenzotriazole
THF	Tetrahydrofuran
TFA	Trifluoroacetic acid
TLC	Thin layer chromatography
TBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate

Pmc 2,2,5,7,8-Pentamethylchroman-6-sulfonyl
 Z Benzyloxycarbonyl

^1H and ^{13}C NMR spectra were recorded using either a Bruker AC 250E spectrometer at 250.1 and 62.9 MHz respectively, or on a Bruker AMX500 at 500.13 and 125.7 MHz respectively. Elemental microanalyses were performed by Medac Ltd (Department of Chemistry, Brunel University, Uxbridge, Middlesex UB8 3PH).

Example 1

4-(O-Methylhydroxylamine)phenoxyethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh)




4-(Hydroxymethyl)phenoxyethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh) ("Wang" resin) (1.83 g, 0.7 mmol/g loading, 1.28 mmol) was suspended in dry THF (20 cm³) and gently agitated for 30 minutes under a blanket of argon. N-Hydroxyphthalimide (624 mg, 3.82 mmol) and triphenylphosphine (1.005 g, 3.82 mmol) were added and the mixture was agitated until these reagents dissolved. Diethylazodicarboxylate (721 mg, 3.28 mmol) was added by cannula to give a bright red solution and the mixture was shaken. Most of the colour dissipated from the reaction mixture after 45 minutes. After 18 h the resin was collected by filtration, washed successively with THF, DMF, dichloromethane, methanol and finally thoroughly with dichloromethane and then dried *in vacuo*; ν_{max} (KBr) 1726(vs), 1687, 1593 cm⁻¹

The resin from above was suspended in DMF (10 cm³) and hydrazine (~100%)(1.5 cm³) was added. The pale yellow solution was heated at 50°C for 2 h and then gently agitated overnight. The desired resin (4-(O-methylhydroxylamine)phenoxyethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh)) was filtered, and washed with DMF (2 x 15 cm³), CH₂Cl₂ (2 x 15 cm³), methanol (20 cm³) and finally with CH₂Cl₂ (3 x 15 cm³) and then dried. Ir (KBr) showed the absence of any carbonyl containing substituent confirming hydrazinolysis of the phthalimide. Elemental analysis of the resin confirmed the presence of nitrogen and suggested a loading of 0.68 mmol/g: C 89.17, H 7.70, N 0.95%. [Note: The loading obtained varied for different batches of the resin due to variability in the loading of the starting Wang resin, but in each case elemental analysis suggested that quantitative conversion was achieved.]

Example 2

O-Hydroxylamine-2'-chlorotrityl-copoly(styrene-1%-divinylbenzene)-resin (200-400 mesh)




Chloro-2'chlorotrityl-copoly(styrene-1%-divinylbenzene) resin (200-400 mesh)(0.5 g, 1.5 mmol/g, 0.75 mmol) was allowed to swell in dry DMF (3 cm³) for 10 minutes. Diisopropylamine (0.398 cm³, 2.28 mmol) was added to a solution of N-hydroxyphthalimide (0.367g, 2.25 mmol) in DMF (3 cm³) and the subsequent bright red solution added in one portion to the resin suspension and the resulting mixture was then gently agitated for 4 hr under an argon atmosphere. The resin was

collected by filtration and washed thoroughly with DMF ($3 \times 10 \text{ cm}^3$), dichloromethane ($2 \times 10 \text{ cm}^3$), followed by methanol ($3 \times 10 \text{ cm}^3$, 3 cycles), dichloromethane ($2 \times 10 \text{ cm}^3$) and dried. The pale yellow resin had ν_{max} (KBr) $1737(\text{vs}) \text{ cm}^{-1}$.

THF (5 cm^3) was added to the resin which was allowed to swell for 15 minutes. Hydrazine hydrate (4 cm^3) was added to the mixture and the resulting mixture was gently agitated for 3 hr. Following filtration and washing as above the resin was dried. It exhibited no stretch at 1737 cm^{-1} in the i.r spectrum and was stained blue by a 2% solution of bromophenol blue (BPB) in DMF.

Example 3

4-[4-(O-Methylhydroxylamine)-3-methoxyphenoxy]-(N-4-methylbenzhydryl)-butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh)




Following the analogous procedure to that described in Example 1, 4-[4-hydroxymethyl-3-methoxyphenoxy]-(N-4-methylbenzhydryl)butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh) (0.87 mmol/g loading) was treated with 3 equivalents of triphenylphosphine, N-hydroxyphthalimide and diethylazodicarboxylate in THF to give the N-hydroxyphthalimido derivatised resin which on treatment with hydrazine hydrate gave the desired 4-[4-(O-

methylhydroxylamine)-3-methoxyphenoxy]-(N-4-methylbenzhydryl)-butyramide-copoly(styrene-1%-divinylbenzene)-resin.

Example 4


4-(2',4'-Dimethoxyphenyl-O-methylhydroxylamine)-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh)



Following the analogous procedure to that described in Example 1, 4-(2',4'-dimethoxyphenyl-O-methylhydroxy)-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh) (0.54 mmol/g loading) was treated with 3 equivalents of triphenylphosphine, N-hydroxyphthalimide and diethylazodicarboxylate in THF to give the N-hydroxyphthalimido derivatised resin which on treatment with hydrazine hydrate gave the desired 4-(2',4'-dimethoxyphenyl-O-methylhydroxylamine)-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin.

Example 5

4-[4-(1-Aminoxyethyl)-2-methoxy-5-nitrophenoxy]-(N-4-methylbenzhydryl)-butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh)



Methyl 4-(4-acetyl-2-methoxyphenoxy)butyrate (10 g) was added to cold 70% nitric acid (140 cm³) and the resulting orange solution was stirred for 2 h and then poured into a beaker containing ice-water (1 dm³). The slurry warmed with stirring overnight to give a yellow solid which was collected and recrystallised from methanol to give methyl 4-(4-acetyl-2-methoxy-5-nitrophenoxy)butyrate as a pale yellow solid (8.5 g). ¹H NMR (CDCl₃) δ 2.21 (2H, m, *J* = 6.6 Hz, CCH₂C), 2.50 (3H, s, MeCO), 2.57 (2H, t, *J* = 7.1 Hz, CH₂CO), 3.7 (3H, s, OMe), 3.99 (3H, s, OMe), 4.16 (2H, t, *J* = 6.2 Hz, CH₂O), 6.75 (1H, s, aromatic) and 7.62 (1H, s, aromatic).

Methyl 4-(4-acetyl-2-methoxy-5-nitrophenoxy)butyrate (5.06 g) in methanol (300 cm³) was cooled with stirring at 0°C under argon. Sodium borohydride (640 mg) was added portionwise and the resulting solution stirred at 40°C overnight. Further sodium borohydride (500 mg) was added as TLC analysis indicated that starting material remained. After a further 3 h TLC analysis indicated that the reaction was complete. The reaction was cooled, acidified with dilute (1M) hydrochloric acid and evaporated to dryness. The residue was dissolved in ethyl acetate (250 cm³) washed with dilute aqueous acid and brine, dried over magnesium sulphate and filtered. The solvent was evaporated to leave a yellow oil which solidified on standing. After recrystallisation from methanol / diethyl ether, methyl 4-[4-(1-hydroxyethyl)-2-methoxy-5-nitrophenoxy]butyrate was obtained as yellow needles (5.04 g, ~ 100%). ¹H NMR (CDCl₃) δ 1.52 (3H, d, *J* = 8.3 Hz, MeCHO), 2.18 (2H, m, CCH₂C), 2.53 (2H, t, *J* = 6.5 Hz, CH₂CO), 3.71 (3H, s, OMe), 3.95 (3H, s, OMe).

4.13 (2H, t, $J = 7.3$ Hz, CH_2O), 5.52 (1H, q, $J = 8.2$ Hz, CHOH), 7.29 (1H, s, aromatic) and 7.57 (1H, s, aromatic).

Methyl 4-[4-(1-hydroxyethyl)-2-methoxy-5-nitrophenoxy]butyrate (5.04 g) from above was stirred with lithium hydroxide monohydrate in methanol using conventional protocols. 4-[4-(1-Hydroxyethyl)-2-methoxy-5-nitrophenoxy]butyric acid was obtained as an amorphous yellow solid (4.0 g, 83%). ^1H NMR (CDCl_3) δ 1.34 (3H, d, $J = 11$ Hz, MeCOH), 2.03 (2H, m, CCH_2C), 2.38 (2H, t, $J = 7.3$ Hz, CH_2CO), 3.81 (3H, s, OMe), 3.96 (2H, t, $J = 7.4$ Hz, CH_2O), 4.13 (2H, brs, OH), 5.38 (1H, q, $J = 10.8$ Hz, CHOH), 7.19 (1H, s, aromatic, and 7.43 (1H, s, aromatic).

4-Methylbenzhydrylamine-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh) (2.0 g, 0.7 mmol/g loading, 1.4 mmol) was placed in DMF (10 cm^3). 4-[4-(1-Hydroxyethyl)-2-methoxy-5-nitrophenoxy]butyric acid (1.51 g, 4.8 mmol), HOBt (734 mg, 4.8 mmol) and TBTU (1.54 g, 4.8 mmol) were added and the mixture agitated. N-Methylmorpholine (0.88 cm^3 , 8 mmol) was added and the resulting mixture was gently agitated for 18 h in a flask protected from light by aluminium foil. The bright yellow resin (2.45 g) was collected by filtration, thoroughly washed as described in Example 1 and dried *in vacuo*. The Kaiser test was negative which contrasted with that of the unsubstituted resin. ν_{max} (KBr) 1672 (s) cm^{-1} .


The above resin (2.41 g) was placed in a small conical flask under argon and THF (25 cm^3) was added. The resin was allowed to swell for 20 minutes and triphenylphosphine (1.67 g, 6.36 mmol) and N-hydroxyphthalimide (1.038 g, 6.36 mmol) were added. The mixture was shaken together in the dark to dissolve the reagents and then cooled to 0° . Diethylazodicarboxylate (1.0 cm^3 , 6.36 mmol) was added and the solution turned red. After shaking the reaction at room temperature for 48 h the resin was then collected by vacuum filtration, washed as described in Example 1 and dried *in vacuo*. ν_{max} (KBr) 1792, 1734 (vs) and 1669 (s) cm^{-1} .

Hydrazine (2.2 cm^3) was added to a suspension of this resin (2.7 g) in DMF (15 cm^3)

under argon. The mixture was shaken at room temperature in the dark overnight and the resin (4-[4-(1-aminooxyethyl)-2-methoxy-5-nitrophenoxy]-(N-4-methylbenzhydryl)-butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh)) was then collected, washed and dried as described in Example 1. It showed no phthalimide carbonyl stretch in the IR spectrum. ν_{\max} (KBr) 1673 (s) cm^{-1} .

Example A

Use of the resin from Example 1 in the preparation of a peptide hydroxamate derivative: Benzyloxycarbonyl-L-prolyl-L-leucyl-L-alanylhydroxamic acid



The resin (4-(O-methylhydroxylamine)phenoxyethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh)) as prepared in Example 1 (200 mg, 0.14 mmol) was suspended in DMF (6 cm^3). Fmoc-Alanine (217 mg, 0.7 mmol), HOBt (107 mg, 0.7 mmol), pyBOP (364 mg, 0.7 mmol) and N-methyl morpholine (0.123 cm^3 , 1.12 mmol) were added and the suspension was agitated for 4 h. The resin was filtered, washed with DMF and then suspended and agitated in a 20% solution of piperidine in DMF for 40 minutes. The solution was drained from the resin which was washed with DMF, dichloromethane, methanol, dichloromethane and finally resuspended in DMF (6 cm^3). Following the same protocols described above Fmoc-leucine and benzyloxycarbonylproline were then successively coupled to the resin.

On completion the resin was washed thoroughly and dried *in vacuo* overnight. The resin was treated with 95% aqueous TFA (10 cm^3) for 75 minutes. The TFA solution

was collected by filtration, the resin washed with more TFA (~2 cm³) and the combined filtrates evaporated to leave a gum which was triturated with ether to give a white solid. TLC of the white solid indicated a single compound which gave a positive ferric chloride test indicative of a hydroxamic acid. The solid was collected and dried to give the desired benzyloxycarbonyl-L-prolyl-L-leucyl-L-alanyl-hydroxamic acid (54 mg, ~90%); ¹H NMR (CDCl₃) δ 7.35 (5H, m, aromatic), 5.15 (1H, J = 7 Hz, CH₂H₆ O), 5.13 (1H, d, J = 7 Hz, CH₂H₆ O), 4.48 (1H, m), 4.33 (1H, m), 4.23 (1H, m), 3.52 (2H, m, pro), 1.9-2.2 (3H, m, pro), 1.48-1.78 (2H, m), 1.48 (3H, d, J = 8.2 Hz, ala), 1.25 (2H, m, CH₂ leu), 0.86 (6H, m, CH₃-leu); ¹³C NMR (CDCl₃) δ 173.5, 172.2, 169.3, 156.4, 136.0, 128.6, 128.4, 127.9, 67.8, 61.5, 53.2, 47.4, 47.3, 39.7, 29.6, 25.1, 24.7, 22.6, 21.4, 17.0.

Example B

Use of the resin from Example 1 in the preparation of a library of 'left hand side' inhibitors of matrix metalloproteinases

Z-AA₃-AA₂-AA₁-NHOH

1g of the resin (4-(O-methylhydroxylamine)phenoxymethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh)) of Example 1 was placed in each of 10 sealable polypropylene tubes (BioRad) and swelled in DMF (1 cm³). 2.5 mmol (4 equivalents) of one of the Fmoc-amino acids, alanine, D-alanine, phenylalanine, isoleucine, leucine, D-leucine, norleucine, methionine, valine, norvaline, was placed in a tube followed by HOBt (3.85 cm³ of a 0.65 M solution in DMF). Diisopropylcarbodiimide (0.4 cm³) was added to each tube which were then sealed and gently agitated for 18 h when they were drained of solvent and washed thoroughly with DMF. A solution of 30% piperidine in DMF (6 cm³) was added to each tube, the resin was mixed by shaking and then allowed to stand in the solution for 40 minutes, the solvent was drained and the resin washed successively with DMF, methanol and dichloromethane as described above in Example 1 and dried.

The resin from each individual tube was then portioned into 10 tubes (~100 mg/tube) and a 1% solution of bromophenol blue in DMF (1 cm³) added to swell and stain the resin.

0.25 mmol of one of the Fmoc-amino acids alanine, D-alanine, arginine (N^ε-Pmc), aspartic acid- β -tert-butyl ester, leucine, proline, norleucine, glutamine- γ -trityl, thioproline, ϵ -Boc-lysine was added to one of the tubes derived from the sub-division of the resin from the original ten tubes. HOBt (0.38 cm³ of a 0.65 M solution in DMF) was added to every tube followed by diisopropylcarbodiimide (39 mm³). The tubes were then sealed and agitated overnight. At this time the colour had discharged from almost every tube. A further 0.05 mmol portion each of the Fmoc-amino acid and diisopropylcarbodiimide was added to those tubes which still showed a blue-green colouring; this was discharged within 30 minutes of the addition of the extra reagent. The resin from these tubes was then pooled back into the tube from where it had originated, washed with DMF and then treated with 30% piperidine in DMF for 40 minutes and washed thoroughly as above.

The resin from each of these ten tubes was then redivided into 5 tubes (~200 mg/tube) and allowed to swell in a 1% solution of bromophenol blue in DMF (1 cm³). 5 mmol of one of the Z-amino acids, alanine, phenylalanine, proline, leucine, aspartic acid- β -tert-butyl ester, was added to one tube from each sub-division followed by HOBt (5 cm³ of a 1.0 M solution in DMF) and by diisopropylcarbodiimide (0.78 cm³). The tubes were gently agitated for 5 h by when the blue colour had been discharged in all tubes. The resin was then repooled into the original tubes as above and thoroughly washed with DMF, dichloromethane and methanol as described previously. The resin, maintained in the 10 tubes, was then dried *in vacuo* overnight.


5 cm³ of a solution of 5% phenol, 5% water, 2% triisopropylsilane in TFA was added to the resin in each tube and periodically shaken over 60 minutes. The TFA solution was collected by filtration and the resin washed with a further 2 cm³ of TFA. The

washings and filtrate from each individual tube were combined and the solvent evaporated to leave a residue which was triturated with cold diisopropylether/hexane to give white solid precipitates. After removal of most of the organic solvent by decanting the precipitates were then lyophilised from 50% aqueous acetonitrile to give the desired 10 mixtures of 50 tripeptide hydroxamic acids.

Each of the ten mixtures of peptidyl hydroxamic acids was assayed for activity as inhibitors of metalloproteinase enzymes. Those pools showing the greatest inhibitory activity were then subjected to an iterative deconvolution procedure to identify individual active inhibitors.

Example C

Use of the resin from Example 1 in the preparation of a sulfonamide hydroxamate derivative: 2-(N-Decyl-2-acetamido-4-methylthiazol-5-ylsulfonamido)-acetohydroxamic acid




Bromoacetic acid (167 mg, 1.2 mmol) and diisopropylcarbodiimide (94 mm³, 0.6 mmol) were added to a suspension of the resin (4-(O-methylhydroxylamine)phenoxymethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh)) prepared according to Example 1 (250 mg, 0.1 mmol) in DMF (6 cm³). The mixture was agitated for 90 minutes and then the solution was drained from the resin which was washed and dried as described above in Example 1. The resin was soaked in DMSO (2 cm³) and a solution of n-decylamine in DMSO (5 cm³ of a 2M solution, 10 mmol) was added and the mixture shaken for 4 h before the resin was

drained and washed thoroughly as before. The resin was then suspended in DMF and 2-acetamido-4-methyl-5-thiazolesulfonylchloride (102 mg, 0.4 mmol) was added. The mixture was heated at 60° for 8 h and then the resin was drained, washed and dried as described previously. Cleavage from the resin as in Example A gave a white solid which was purified by column chromatography (10% methanol - dichloromethane; silica) to give the desired 2-(N-decyl-2-acetamido-4-methylthiazol-5-ylsulfonamido)-acetohydroxamic acid (23 mg, 51%) as a white solid; ¹H NMR (CDCl₃) δ 3.83 (2H, m), 3.05 (2H, m), 2.52 (3H, s, Me), 2.35 (3H, s, NHCOMe), 1.69 (2H, m), 1.15-1.25 (14H, m, alkyl), 0.87 (3H, t, J = 7.2Hz, CH₃-alkyl).

Example D

Use of the resin from Example 1 in the preparation of an array of sulfonamide hydroxamate inhibitors of matrix metalloproteinases



Sulfonamide hydroxamic acid derivatives were prepared as a combinatorial array of individual compounds in purity sufficient for fast throughput assays. The resin (4-(O-methylhydroxylamine)phenoxyethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh)) was loaded with bromoacetic acid as described in Example C.

Elemental analysis of this different batch of resin confirmed that coupling had taken place: C 83.34, H 7.04, N 0.86, Br 5.40% and suggested a loading of 0.65 mmol. 1 g of this resin was placed in each of 36 plastic Econopac tubes (BioRad) and swelled in DMF (5 cm³). The excess DMF was drained off and the tubes grouped into nine groups of four tubes. A solution (5 cm³ of a 2M solution in either DMF or DMSO) of one of the amines, benzylamine, 4-methylbenzylamine, 4-methoxybenzylamine, 4-

fluorobenzylamine, 4-chlorobenzylamine, pyridin-3-ylmethylamine, thien-2-ylmethylamine, furan-2-ylmethylamine and 3,4-dioxymethylenebenzylamine was each added to each of four tubes (within one of the nine groups) which were then sealed and mixed thoroughly for 2 h. The solvent was removed, the resin washed as in Example C and reswelled in DMF. The tubes were regrouped into four groups each of nine tubes. Following a protocol analogous to that used in Example C, a DMF solution of 10 equivalents of one of the sulfonyl chlorides, 3-chloropropylsulfonyl chloride, hexylsulfonyl chloride, octylsulfonyl chloride and decylsulfonyl chloride was added to each of the nine tubes (within one of the four groups) so that all 36 combinations of sulfonamides were prepared. After removal of the solvent by filtration, thorough washing with DMF, methanol and dichloromethane as above, and drying under vacuum for several hours the products were cleaved from the resin as in Example C directly into plastic centrifuge tubes. The TFA was removed by centrifugal evaporation and the resulting hydroxamic acids lyophilised from aqueous acetonitrile. All samples gave positive ferric chloride spots on TLC analysis {10% methanol in DCM/silica} (Table). All 36 samples were submitted for assay as inhibitors of metalloproteinases from which active compounds were identified.


Table: TLC {10% methanol in DCM/silica} R_f values for the compounds of Example D

Ra	Rb			
	$-(CH_2)_3Cl$	$-C_6H_{13}$	$-C_8H_{17}$	$-C_{10}H_{21}$
C_6H_5-	0.25	0.2	0.45	0.45
4-Me C_6H_4-	0.35	0.4	0.4	0.4
4-MeOC $_6H_4-$	0.5	0.35	0.5	0.4
4-FC $_6H_4-$	0.3	0.3	0.5	0.5
4-ClC $_6H_4-$	0.15	0.2	0.4	0.4
pyridin-3-yl	0.0	0.05	0.0	0.05
thien-2-yl	0.1	0.35	0.65	0.6

furan-2-yl	0.15	0.3	0.3	0.4
3,4-(CH ₂)O ₂ C ₆ H ₃ -	0.3	0.4	0.5	0.45

Example E

Use of the resin from Example 2 in the preparation of a peptide hydroxamate derivative: Benzyloxycarbonyl-L-phenylalanyl-L-alanylhydroxamic acid




FMOC-Alanine (187 mg, 0.6 mmol), HOBt (92 mg, 0.6 mmol) and TBTU (194 mg, 0.6 mmol) were dissolved together in DMF (5 cm³). Diisopropylethylamine (0.183cm³, 1.05 mmol) was added and the resulting solution added to pre-swelled (in DMF) modified resin from Example 2 (O-hydroxylamine-2'-chlorotrityl-copoly(styrene-1%-divinylbenzene)-resin (200-400 mesh)) (100 mg, 0.15 mmol) and the mixture was then shaken gently together for 12 h. The resulting resin was then filtered, washed thoroughly with DMF and treated with 20% piperidine in DMF (7 cm³) for 30 minutes. After removal of the solvent and thorough washing as described in Example 1 the resin was resuspended in DMF (1 cm³). A solution of Z-phenylalanine (180 mg, 0.6 mmol), HOBt (92 mg, 0.6 mmol) and diisopropylcarbodiimide (94 mm³, 0.6mmol) in DMF (5 cm³) was added and the resin suspension was gently agitated overnight by which time the initially blue staining of BPB had been discharged. The resin was then drained and washed as described in Example 1 and dried *in vacuo*. It had ν_{max} (KBr) 1668 (s) cm⁻¹.

The resin from above was suspended in a 20% solution of TFA in dichloromethane, containing 1% (v/v) of triethylsilane, under argon (5 cm³) for 45 minutes. The

solution was removed from the resin by filtration, the resin washed with a further portion (1 cm³) of the cleavage mixture, the filtrate and washings combined and evaporated to leave benzyloxycarbonyl-L-phenylalanyl-L-alanylhydroxamic acid as a white solid. TLC analysis indicated a single compound, R_f 0.45 (10% methanol in dichloromethane) which gave a positive ferric chloride test. ¹H NMR (CDCl₃) δ 1.36 (3H, d, J = 9 Hz, Me alanine), 3.12 (2H, m, CH₂Ph), 4.14 (1H, m, CH), 4.4 (1H, m, CH), 5.13 (2H, brs, OCH₂Ph), 7.71-7.33 (~10H, m, aromatic) 7.8 (1H, d, NH).

Example F

Use of the resin from Example 3 in the preparation of a sulfonamide hydroxamate derivative: 2-(N-Benzyl-benzylsulfonamido)-acetohydroxamic acid



The resin of Example 3 (4-[4-(O-methylhydroxylamine)-3-methoxyphenoxy]-(N-4-methylbenzhydryl)butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh)) (0.5 g) was suspended in DMF (3 cm³). Diisopropylcarbodiimide (0.3 cm³) was added to a solution of bromoacetic acid (670 mg) in DMF (3 cm³) and after 3 minutes the resulting solution was added to the resin suspension and gently mixed together for 2 h. The resin was collected by filtration and washed thoroughly with DMF, methanol, methanol-dichloromethane, dichloromethane and ether. It was then dried *in vacuo* at 45°C for 18 h. Elemental analysis of the resin suggested a loading of 0.66 mmol/g: C; 79.11; H, 6.78; N, 2.08; Br, 5.25%.

The resin from above swelled in DMF (3 ml) was treated with (0.33 cm³, 3 mmol)

benzylamine and gently agitated at room temperature overnight. The solvent was removed by filtration and the resin washed as described above in Example 1 and dried. The resin was swelled in DMF (3ml), and phenylmethanesulfonyl chloride (0.8 g, 4 mmol) was added and the mixture was gently agitated for 2 h at room temperature. The resin was filtered and washed as described above in Example 1 and dried. A solution of 1% TFA in DCM (3 ml) was added to the resin at room temperature and the mixture gently agitated for 1 h. The resin was filtered and washed with DCM, methanol and DCM, the filtrates combined and evaporated under reduced pressure to give crude 2-(N-benzyl-benzylsulfonamido)-acetohydroxamic acid (0.39 g): TLC ferric chloride positive spot R_f 0.15 {10% methanol in DCM/silica}.

Example G

Use of the resin from Example 4 in the preparation of a sulfonamide hydroxamate derivative: 2-(N-Benzyl-benzylsulfonamido)-acetohydroxamic acid

This resin of Example 4 (4-(2',4'-dimethoxyphenyl-O-methylhydroxylamine)-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh)) (0.25 g) was suspended in DMF (3 cm³). Diisopropylcarbodiimide (0.24 cm³) was added to a solution of bromoacetic acid (420 mg) in DMF (3 cm³) and after 3 minutes the resulting solution was added to the resin suspension and gently mixed together for 2 h. The resin was collected by filtration and washed thoroughly with DMF, methanol, methanol-dichloromethane, dichloromethane and ether. It was then dried *in vacuo* at 45°C for 18 h. Elemental analysis of the resin suggested a loading of 0.58 mmol/g: C, 84.31; H, 7.28; N, 0.82; Br, 4.65%.

Following the procedure of Example F the above resin was treated in turn with benzylamine and phenylmethanesulfonyl chloride. A solution of 10% acetic acid in DCM (3 ml) was added to the resin at room temperature and the mixture gently agitated for 2 h. The resin was filtered and washed with DCM, methanol and DCM, the filtrates combined and evaporated under reduced pressure to give crude 2-(N-

benzyl-benzylsulfonamido)-acetohydroxamic acid (0.22 g).


Example H

Use of the resin from Example 5 in the preparation of a peptide hydroxamate derivative: Benzyloxycarbonyl-L-prolyl-L-leucyl-L-alanylhydroxamic acid


Using procedures analogous to those described in Example A, the tripeptide Z-L-prolyl-L-leucyl-L-alanyl was synthesised attached to the resin of Example 5 (4-[4-(1-aminooxyethyl)-2-methoxy-5-nitrophenoxy]-(N-4-methylbenzhydryl)-butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh)). Following drying, the elaborated resin (50 mg) in acetonitrile (1.5 cm³) was blanketed with argon in a stoppered reaction vial. The vial was placed 5 cm away from a 365 nm light source and irradiated for 15 h. TLC of the resulting solution indicated a ferric chloride positive component with R_F equivalent to that of the tripeptide hydroxamic acid of Example A. HPLC of the solution identified the desired tripeptide as the major product of the reaction along with an unidentified non-hydroxamic acid component.

Claims

1. A solid phase reaction component comprising a solid substrate, substantially insoluble in aqueous or organic reaction media, carrying a plurality of covalently bound hydroxylamine or protected hydroxylamine groups of formula (A) or (B)



(A)




(B)

wherein P_1 is hydrogen or an amino protecting group and P_2 is hydrogen or a hydroxyl protecting group, and the bond designated (a) is one which covalently links the group (A) or (B) to the residue of the solid substrate and is cleavable under acid conditions or by photolysis.

2. A solid phase reaction component as claimed in claim 1 wherein the solid substrate comprises a base substrate carrying linker groups which indirectly link the hydroxylamine group (A) or (B) to the base substrate.

3. A solid phase reaction component as claimed in claim 2 wherein the base substrate carries a plurality of groups of formula (IV):



wherein


R_{10} represents hydrogen or (phenyl) CH_2 -, optionally substituted in the phenyl ring by one or more substituents selected from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, nitrile or NO_2 ;

R_{11} represents hydrogen, C_1 - C_6 alkyl, or phenyl optionally substituted by one or more substituents selected from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, nitrile or NO_2 ;

R_{12} and R_{12A} independently represent hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, nitrile or NO_2 ;

R_{13} represents a group $-(X^1)_q-Y$ - wherein q is 0 or 1, X^1 represents $-\text{C}(=\text{O})-$, $-\text{CH}_2-$, $-\text{CH}_2\text{C}(=\text{O})-$, $-\text{O}(\text{CH}_2)_n\text{C}(=\text{O})-$, $-\text{O}(\text{CH}_2)_n\text{C}(=\text{O})-(A^1)_m-$, or $-\text{O}(\text{CH}_2)_n\text{C}(=\text{O})-(A^1)_m-B^1-$, wherein n is an integer from 1 to 6, m is 0 or 1, A^1 represents $-\text{O}-\text{CH}(R^1)-\text{NH}-$ wherein R^1 is the side chain of a natural or unnatural alpha amino acid, B^1 represents a spacer group $-\text{NH}(\text{CH}_2)_p-$ wherein p is 0 or an integer from 1 to 6, and Y represents $-\text{O}-$ or $-\text{NH}-$.

4. A solid phase reaction component as claimed in claim 2 wherein the base substrate carries a plurality of groups of formula (IVA):



wherein R_{10} , R_{11} , R_{12} and R_{12A} are as defined in claim 3, R_{11A} is as defined for R_{11} in claim 3, and R_{13A} is a bond or is as defined for R_{13} in claim 3.

5. A solid phase reaction component as claimed in claim 3 or claim 4 wherein R_{10} is hydrogen, 4-methoxybenzyl or 2,4-dimethoxybenzyl.



6. A solid phase reaction component as claimed in any one of claims 3 to 5 wherein R_{11} is hydrogen, methyl, phenyl, 4-methylphenyl, 4-methoxyphenyl or 2,4-dimethoxyphenyl.
7. A solid phase reaction component as claimed in any one of claims 3 to 6 wherein R_{12} is hydrogen or methoxy.
8. A solid phase reaction component as claimed in any one of claims 3 to 7 wherein R_{12A} is hydrogen or methoxy.
9. A solid phase reaction component as claimed in claim 3 or any or claims 4 to 8 as dependent on claim 3, wherein R_{13} together with the base substrate to which it is attached is, oxymethyl-copoly(styrene-1% divinylbenzene)resin, oxymethyl-copoly(styrene-2% divinylbenzene)resin, oxyacetomidomethylpolyethyleneglycol-copoly(styrene-1% divinylbenzene)resin or oxyacetomidomethylpolyethyleneglycol-copoly(styrene-2% divinylbenzene)resin.
10. A solid phase reaction component as claimed in claim 1 selected from
 - 4-(O-Methylhydroxylamine)phenoxyethyl-copoly(styrene-1% divinylbenzene)-resin (100-200 mesh),
 - 4-[4-(O-Methylhydroxylamine)-3-methoxyphenoxy]-(N-4-methylbenzhydryl)butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh),
 - 4-(2',4'-Dimethoxyphenyl-O-methylhydroxylamine)-phenoxyethyl-copoly(styrene-1%-divinylbenzene)-resin (100-200 mesh),
 - 4-[4-(1-Aminooxyethyl)-2-methoxy-5-nitrophenoxy]-(N-4-methylbenzhydryl)butyramide-copoly(styrene-1%-divinylbenzene)-resin (100-

200 mesh), or

O-Hydroxylamine-2'-chlorotriyl-copoly(styrene-1%-divinylbenzene)-resin
(200-400 mesh)

11. A process for the preparation of a desired synthetic product whose structure is characterised by the presence of a covalently bonded hydroxamic acid group - CONHOH, which process comprises the steps of:

(i) forming a mixture of a liquid reaction medium and a solid phase reaction component which is substantially insoluble in the said liquid reaction medium and carries a plurality of moieties of formula (A1) or (B1)



in which formulae X represents the residual, non-hydroxamate, partial structure of the desired product, P₁ represents hydrogen or an amino-protecting group, P₂ represents hydrogen or a hydroxyl protecting group, and the bond designated (a) is one which covalently links the moieties (A1) or (B1) to the residue of the solid substrate and is cleavable under acid conditions or by photolysis; and

(ii) in the resultant mixture, cleaving the said bond designated (a) and, if P₁ or P₂ as the case may be is not hydrogen, removing that protecting group P₁ or P₂ before, after or during cleavage of bond (a); and

(iii) separating the resultant liquid reaction phase from the resultant reaction solids and recovering the desired product from the separated liquid reaction phase.

12. A process as claimed in claim 11 wherein the solid phase reaction component which carries a plurality of moieties of formula (A1) or (B1) is in the form of a finely divided solid, or a web, membrane or open pore matrix.
13. A process as claimed in claim 11 or claim 12 wherein in step (i) the solid phase reaction component which carries a plurality of moieties of formula (A1) or (B1) is derived by chemical modification from a solid phase reaction component as claimed in any one of claims 1 to 10.

INTERNATIONAL SEARCH REPORT

In tional Application No
PCT/GB 96/00428

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C08F8/30 C12N11/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C08F C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 561 722 (BIO MERIEUX) 22 September 1993 see claims 1-18	1
A	EP,A,0 216 499 (COURTAULDS PLC) 1 April 1987 see page 5, line 22 - page 6, line 14; claims 1-11	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

24 May 1996

Date of mailing of the international search report

12.06.1996

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Permentier, W

INTERNATIONAL SEARCH REPORT

In International Application No
PCT/GB 96/00428

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMICAL ABSTRACTS, vol. 77, no. 24, 11 December 1972 Columbus, Ohio, US; abstract no. 153114e, CHERNYSHEV, V. P.: "MALEIC ANHYDRIDE COPOLYMER" page 39; column 2; XP002003869 see abstract & SU,A,342 865 (INSTITUTE OF FINE CHEMICAL TECHNOLOGY, MOSCOW) 22 June 1972 ---	1
A	EP,A,0 514 648 (AMERICAN CYANAMID COMPANY) 25 November 1992 see claims 1-10 ---	1
A	US,A,4 868 248 (J. W. SPARAPANY) 19 September 1989 see the whole document ---	1
A	US,A,4 461 876 (S. LIEBERMAN) 24 July 1984 see claims 1-13 ---	1
A	POLYMER BULLETIN, vol. 32, no. 3, 1994, pages 273-279, XP002003868 TAEK SEUNG LEE: "SYNTHESIS OF POROUS POLY(HYDROXAMIC ACID) FROM POLY(ETHYL ACRYLATE-CO-DIVINYLBENZENE)" see page 273 - page 279 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

In tional Application No
PCT/GB 96/00428

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-561722	22-09-93	FR-A- 2688788 US-A- 5489653 US-A- 5439972	24-09-93 06-02-96 08-08-95
EP-A-216499	01-04-87	NONE	
EP-A-514648	25-11-92	AU-B- 651885 AU-B- 1710092 CA-A- 2069169 JP-A- 5194632	04-08-94 26-11-92 24-11-92 03-08-93
US-A-4868248	19-09-89	NONE	
US-A-4461876	24-07-84	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.